



Chapter 13 - Chemical Equilibrium



The Equilibrium State

- Figure 13.1



The Equilibrium State

- Not all chemical reactions go to completion.
- When you hear “equilibrium”, what do you think of?
- Example: weather patterns: ocean water evaporates at the same rate that it rains. They are in equilibrium.

The Equilibrium State

- We commonly use concentrations to describe equilibrium.
 - ♦ $[] = M = \text{moles/L}$
 - ♦ At equilibrium, the concentrations of substances remain constant.
 - ♦ The rate of the forward reaction is the same as the rate of the reverse reaction.
- Usually one side of the equation is favored (reactants or products).

The Equilibrium State

- Equilibrium: The state reached when the **concentrations** of reactants and products remain constant over time.
- Most reactions are reversible.
 - ♦ $A + B \leftrightarrow C + D$
 - ♦ A and B react to make C and D
 - ♦ But C and D can also react to make A and B

The Equilibrium Constant, K_c




- The Equilibrium Constant, K_c , tells us which side of the reaction is favored.
- $2 \text{H}_2 (\text{g}) + \text{O}_2 (\text{g}) \leftrightarrow 2 \text{H}_2\text{O} (\text{g})$
- $K_c = \frac{[\text{H}_2\text{O}]^2}{[\text{H}_2]^2[\text{O}_2]}$
 - ♦ K_c is the equilibrium constant, a value
 - ♦ The fraction is the equilibrium constant expression
- K_c is constant at a particular T, K_c is unitless.

Equilibrium equation: $K_c = \frac{[\text{C}]^c[\text{D}]^d}{[\text{A}]^a[\text{B}]^b}$

↙ Equilibrium constant ↘ Equilibrium constant expression

The Equilibrium Constant, K_c

- Write the equilibrium constant expressions (K_c) for the following equations. (Note: Expressions don't include solids or liquids!)



★ $K_c = \frac{[\text{CH}_4][\text{H}_2\text{O}]}{[\text{CO}][\text{H}_2]^3}$

Ⓢ $K_c = \frac{[\text{N}_2][\text{H}_2]^3}{[\text{NH}_3]^2}$

Ⓢ $K_c = \frac{[\text{SO}_3]^2}{[\text{SO}_2]^2[\text{O}_2]}$

Calculating K_c

- $2\text{NH}_3\text{(g)} \leftrightarrow \text{N}_2\text{(g)} + 3\text{H}_2\text{(g)}$
- At 500 K, the following concentrations were measured: $[\text{N}_2] = 3.0 \times 10^{-2} \text{ M}$, $[\text{H}_2] = 3.7 \times 10^{-2} \text{ M}$, $[\text{NH}_3] = 1.6 \times 10^{-2} \text{ M}$. What is K_c ?
- Are you writing an expression or solving for a number?
- Problems 13.1 - 13.4

The Equilibrium Constant, K_c

- $2 \text{H}_2\text{(g)} + \text{O}_2\text{(g)} \leftrightarrow 2 \text{H}_2\text{O(g)}$
- If $K_c = 2.4 \times 10^{47}$ (e.g., a large number) what does this tell you about the **equilibrium** of the reaction?
 - A. There are more reactants than products
 - B. There are more products than reactants
 - C. More information is needed.
- How can you figure this out?

Group Quiz #3


- $\text{N}_2\text{O}_4\text{(g)} \leftrightarrow 2\text{NO}_2\text{(g)}$
- At 100°C, the following concentrations are measured: $[\text{N}_2\text{O}_4] = 0.0172 \text{ M}$, $[\text{NO}_2] = 0.00140 \text{ M}$. What is K_c at this temperature?

The Equilibrium Constant, K_c

- $2 \text{HBr(g)} \leftrightarrow \text{H}_2\text{(g)} + \text{Br}_2\text{(g)}$
- $K_c = 2 \times 10^{-19}$ (e.g., a small number), what does this tell you about the **equilibrium** of the reaction?
 - A. There are more reactants than products
 - B. There are more products than reactants
 - C. More information is needed.

Interpreting the Equil. Constant

- If $K_c > 10^3$, products are favored over reactants; reaction goes nearly to completion.
- If $K_c < 10^{-3}$, reactants are favored over products; reaction hardly proceeds at all.
- If K_c is in the range $10^{-3} - 10^3$, appreciable concentration of reactants and products are present.
- Problem 13.8



K_c versus K_p

- Can measure gas pressures instead of molar concentrations. Pressure is directly proportional to concentration.
- $PV = nRT$ $\Leftrightarrow P = (n/V) RT \Leftrightarrow n/V = M$
- $2\text{NH}_3(\text{g}) \leftrightarrow \text{N}_2(\text{g}) + 3\text{H}_2(\text{g})$
 - ♦ $K_c = [\text{N}_2][\text{H}_2]^3 / [\text{NH}_3]^2$
 - ♦ $K_p = (P_{\text{N}_2})(P_{\text{H}_2})^3 / (P_{\text{NH}_3})^2$
- $K_p = K_c(RT)^{\Delta n}$
 - ♦ $\Delta n = \text{moles product} - \text{moles reactant}$
 - ♦ $R = 0.08206 \text{ L}\cdot\text{atm} / \text{mol}\cdot\text{K}$

Heterogeneous Equilibria

- Write K_c and K_p for the following equations:
- $\text{CaCO}_3(\text{s}) \leftrightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
- $2\text{Cu}_2\text{S}(\text{s}) + 3\text{O}_2(\text{g}) \leftrightarrow 2\text{Cu}_2\text{O}(\text{s}) + 2\text{SO}_2(\text{g})$
- $\text{Hg}(\text{l}) + \text{Hg}^{2+}(\text{aq}) \leftrightarrow \text{Hg}_2^{2+}(\text{aq})$

Worked Example 13.5

- $\text{CH}_4(\text{g}) + 2\text{H}_2\text{S}(\text{g}) \leftrightarrow \text{CS}_2(\text{g}) + 4\text{H}_2(\text{g})$
- At 1000 K, CH_4 is 0.20 atm, H_2S is 0.25 atm, CS_2 is 0.52 atm, and H_2 is 0.10 atm. What is K_p ?
- Ans: 4.2×10^{-3}

Using the Equil. Constant

- $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{H}_2\text{O}(\text{g})$
 - ♦ $K_c = 2.4 \times 10^{47}$ at 500K
 - ♦ What is the concentration of H_2 at equilibrium if $[\text{H}_2\text{O}] = 1.0 \text{ M}$ and $[\text{O}_2] = 1.0 \times 10^{-16} \text{ M}$?
- $3\text{H}_2(\text{g}) + \text{N}_2(\text{g}) \leftrightarrow 2\text{NH}_3(\text{g})$
 - ♦ Calculate K_c : $[\text{H}_2] = 0.104 \text{ M}$, $[\text{N}_2] = 0.554 \text{ M}$, $[\text{NH}_3] = 0.418 \text{ M}$
 - ♦ Calculate P_{NH_3} : $P_{\text{H}_2} = 1.24 \text{ atm}$, $P_{\text{N}_2} = 2.17 \text{ atm}$

Heterogeneous Equilibria

- Homogeneous equil.: all substances are in one phase (all gas, all solid, etc.)
- Heterogeneous equil.: substances in 2 or more different phases
- $\text{CaCO}_3(\text{s}) \leftrightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
- Solids and liquids have constant concentrations. If you increase the amount of CaCO_3 , you also increase its volume.

Reaction Quotient, Q

- Q is similar to K but may not be at equilibrium. Calculating Q tells us which direction a reaction must go to reach equilibrium.
- $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \leftrightarrow 2\text{HI}(\text{g})$ $K_c = 57.0$ at 700 K
- If $[\text{H}_2] = 0.10 \text{ M}$, $[\text{I}_2] = 0.20 \text{ M}$, and $[\text{HI}] = 0.40 \text{ M}$, is this system at equilibrium?
- What needs to happen to reach equilibrium?

Reaction Quotient, Q


- If $Q_c < K_c$, reaction shifts right (creates more products)
- If $Q_c > K_c$, reaction shifts left (creates more reactants)
- If $Q_c = K_c$, reaction is at equilibrium (no shifting)
- Figure 13.5
- Worked Ex 13.8, Problems 13.9 - 13.10

Calculate Equil. Concentrations

- Given initial concentrations of reactants and K_c , we can calculate equilibrium concentrations.
- Use balanced equation and an ICE table.

$K_c = 2.5 \times 10^{-5}$	$N_2(g)$	$3H_2(g)$	\leftrightarrow	$2NH_3(g)$
Initial	1.00 M	1.00 M		0 M
Change	-x	-3x		+2x
Equilibrium	1.00 - x	1.00 - 3x		2x

Reaction Quotient, Q



Calculate Equil. Concentrations

- $N_2(g) + 3H_2(g) \leftrightarrow 2NH_3(g)$
- Is the equation balanced?
- $[N_2]_i = 1.00\text{ M}$, $[H_2]_i = 1.00\text{ M}$, $[NH_3]_i = 0\text{ M}$
- What are equilibrium concentrations of all species?
- $K_c = 2.5 \times 10^{-5}$
- Ans: $[N_2]_{eq} = 0.998\text{ M}$, $[H_2]_{eq} = 0.993\text{ M}$, $[NH_3]_{eq} = 5.00 \times 10^{-3}\text{ M}$

Direction of Shift

- $CO(g) + 3H_2(g) \leftrightarrow CH_4(g) + H_2O(g)$ $K = 4.0$
2.0 M 1.0 M 0.50 M 0.50 M
- Is this system at equilibrium? In which direction will the reaction proceed to reach equilibrium?
- Group Quiz #4:
- $CO(g) + 3H_2(g) \leftrightarrow CH_4(g) + H_2O(g)$ $K = 4.0$
0.20 M 0.10 M 1.0 M 1.0 M
- Is this system at equilibrium? In which direction will the reaction proceed to reach equilibrium?

PRACTICE!!!

- Worked examples 13.9 - 13.11
- Problems 13.11 - 13.15

Solving for Equil. Conc.

- Assumption that x is small isn't always valid!
- What to do?
- Quadratic formula!!!

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- Gives two values of x; both won't always make sense

Group Quiz #5 Answers

- $K_c = 4.00 = [\text{CO}_2][\text{H}_2] / [\text{CO}][\text{H}_2\text{O}]$
- $4.00 = x^2 / (1.00 - x)(2.00 - x)$
- Assume $x \ll 1.00$, $x \ll 2.00$
- $x = 2.83$
- Assumption invalid; quadratic!
- $3.00x^2 - 12.00x + 8.00 = 0$
 - $[\text{CO}_2]_{\text{eq}} = [\text{H}_2]_{\text{eq}} = x = 0.845 \text{ M}$
 - $[\text{CO}]_{\text{eq}} = 1.00 - x = 0.155 \text{ M}$
 - $[\text{H}_2]_{\text{eq}} = 2.00 - x = 1.155 \text{ M}$
- Check math by plugging in concentrations!

Solving for Equil. Conc.

- $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \leftrightarrow 2\text{HI}(\text{g})$
1.00 M 2.00 M 0 M
- $K_c = 50.5$
- Calculate the equilibrium concentrations of all species.
- Check math by plugging concentrations in to equilibrium expression.

Factors Affecting Equil.

- Now we look at what factors can cause shifts in equilibrium. These can alter expenses in industrial settings where a shift toward creation of more products is often more economical.
 - Concentrations of reactants or products can be changed.
 - Pressure and volume of the system can be changed.
 - Temperature can be changed.
 - Addition of a catalyst.

Group Quiz #5

- $\text{CO}(\text{g}) + \text{H}_2\text{O}(\text{g}) \leftrightarrow \text{CO}_2(\text{g}) + \text{H}_2(\text{g})$
1.00 M 2.00 M 0 M 0 M
- $K_c = 4.00$
- What are the equilibrium concentrations?

Le Chatelier's Principle

- If a stress is applied to a reaction mixture at equilibrium, net reaction occurs in the direction that relieves the stress.
- "Stress" means a change in one of the factors mentioned on the previous screen.

Changes in Concentration

- $A \leftrightarrow B$ $K = 4$
- At equilibrium, $[A] = 5$ and $[B] = 20$
- I remove 5 B's. Is this still at equilibrium? If not, what is Q ? Which way does this system need to shift? Can you figure this out non-mathematically?
- What happens if I add 5 A's?

Changes in Concentration

- $N_2(g) + 3 H_2(g) \leftrightarrow 2 NH_3(g)$
1.50 M 3.00 M 2.00 M
- $K_c = 0.296$ at 700 K
- Calculate equilibrium concentrations

Changes in Concentration


- $N_2(g) + 3 H_2(g) \leftrightarrow 2 NH_3(g)$
- $K_c = 0.296$ at 700 K
- At equilibrium, $[N_2] = 0.50$ M, $[H_2] = 3.00$ M, $[NH_3] = 2.00$ M
- Verify K_c

Changes in P and V

- Increase in volume: more space = more gases allowed. Shift to side with more **moles of gas**.
- Increase in pressure (same as decrease in volume): less volume = less gas allowed. Shift to side with fewer **moles of gas**.
- If same number of moles on both sides, P and V don't affect equilibrium.
- Adding an inert gas also won't change anything.

Adding Reactant, N_2

- What will happen if we add N_2 to the system?
- Figure 13.8



Changes in P and V

- $N_2(g) + 3 H_2(g) \leftrightarrow 2 NH_3(g)$
- There are 4 moles of reactant and 2 moles of product.
- What will happen if we increase pressure?
- What will happen if we increase volume?
- Example 13.13 and Problem 13.17

Changes in Temperature


- Changes in Conc., P, and V just shift equilibrium to maintain a constant K.
- Changes in Temperature usually affects the value of K.
- We can look at heat exchange (enthalpy) of a reaction to predict shift.
- Endothermic: heat is a “reactant”
- Exothermic: heat is a “product”

Le Chatelier's Principle

- Determine how the equilibrium will shift if the following changes are made:
- $2\text{Cl}_2(\text{g}) + 2\text{H}_2\text{O}(\text{g}) \leftrightarrow 4\text{HCl}(\text{g}) + \text{O}_2(\text{g})$
 $\Delta H^\circ = +113 \text{ kJ}$
- Temperature is increased
- Volume is increased
- Pressure is decreased
- HCl is added
- Ne (g) is added
- Cl_2 is added
- H_2O is removed

Changes in Temperature

- $2\text{NO}_2(\text{g}) \leftrightarrow \text{N}_2\text{O}_4(\text{g})$ $\Delta H^\circ = +57.2 \text{ kJ}$
- Endothermic, heat can be imagined to be a reactant, it must be put in to the system.
- Add heat, adds reactant
- Shifts right
- Figure 13.13



Le Chatelier's Principle

- Determine how the equilibrium will shift if the following changes are made:
- $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2\text{H}_2\text{O}(\text{g})$ $\Delta H^\circ = -571 \text{ kJ}$
- Temperature is increased
- Volume is decreased
- Pressure is decreased
- H_2 is added
- O_2 is removed
- H_2O is removed

Adding a Catalyst

- Catalysts speed up a reaction so they reach equilibrium faster. Once at equilibrium, catalysts serve no purpose.
- If a system is already at equilibrium, adding a catalyst won't affect the system at all!